

A non standard approach of spectral theory of self-adjoint operators (generalized Gelfand eigenvectors)

Fatma Karray Meziou
 Université Pierre & Marie Curie.
 meziou@math.jussieu.fr

Abstract

Let $u = \int_{-\infty}^{+\infty} \lambda dE_\lambda$ be a self-adjoint operator in a Hilbert space H . Our purpose is to provide a non-standard description of the spectral family (E_λ) and the generalized Gelfand eigenvectors.

1 The non standard framework

We consider here the nonstandard analysis framework (see [Goldblatt], for a brief overview) and apply it to set universes : Let A be a set of $\in$ -independent elements (i.e. a non empty set such as there is not any chain $x_0 \in x_1 \in \dots \in x_n$ whose ends belong to A) and let us associate to A its superstructure U_A , i.e. the union of the sequence (A_n) defined by $A_0 = A$ and $A_{n+1} = A_n \cup P'(A_n)$, where P' denotes the set of non empty subsets. The couple made of U_A and the restriction of $\in$ to U_A is a standard model of Zermelo theory with atoms and where Zorn's theorem holds.

Note :

1. Zermelo theory is ZF without the substitution axiom (see [Jech]).
2. We need to choose once for all a $\emptyset_A \in A$ that will play the role of the empty set in U_A ; U_A atoms are then the elements of $A \setminus \{\emptyset_A\}$. Such a U_A universe contains all necessary sets to describe the structures (in the sense of N.Bourbaki) over elements of U_A .
3. We may assume that A is chosen such that $\mathbb{C}$ and H are its subsets; even if this requires to replace $\mathbb{C}$ and H by isomorphic copies. $\mathbb{C}$ and H have then their binary operation graphs in U_A . One can associate to U_A a first order language $L(U_A)$ whose relational symbols are $=$ and $\in$ and whose constant set is U_A .

The non-standard method relies on the existence of a function called enlargement defined from U_A to the set class as $a \mapsto^* a$ with the following properties:

- i. $^*a = a$ for each atom,
- ii. if R is a relation of $L(U_A)$ without any free variable and where all quantifiers are limited - i.e. in the form of $\exists x \in t$ or $\forall x \in t$ - and if *R is the relation obtained by replacing every outstanding constant a by its image *a , R and *R relations are either both true or both false (transfer principle),
- iii. if $F \in U_A$ is a filter on a set $u \in U_A$, the intersection of *F sets (which is a filter over *u) is not empty.

In particular, given i and ii, $\mathbb{C}$ could be enlarged into a set $^*\mathbb{C}$ containing $\mathbb{C}$. We will call $^*\mathbb{C}$ elements hypercomplexes and define over $^*\mathbb{C}$ an addition and a multiplication operations, whose graphs are the enlargements of the addition and multiplication graphs over $\mathbb{C}$. For these operations $^*\mathbb{C}$ is an algebraically closed field. The same analysis applies to $\mathbb{R}$ and leads to the subfield $^*\mathbb{R}$ of $^*\mathbb{C}$. $^*\mathbb{R}$ is totally ordered and generates with element i the field $^*\mathbb{C}$ (This allows to consider the real and imaginary parts of a hypercomplex and to introduce the absolute hypervalue of a hypercomplex).

An element $\xi \in ^*\mathbb{R}$ is infinitesimal (resp. limited), if $-1 < n\xi < 1$ for every natural number n (resp. $-n < \xi < n$ for at least one natural number). A hypercomplex is infinitesimal (resp. limited) if its absolute hypervalue is infinitesimal (resp. limited). The subset $^*\mathbb{C}_b$ made of bounded elements is a valuation ring whose residual field could be identified with $\mathbb{C}$: more precisely the image of $z \in ^*\mathbb{C}_b$ is the unique complex number ${}^\circ z$ such that $z - {}^\circ z$ is infinitesimal.

From now on, we consider a fixed universe U and an enlargement *U of U .

2 Hyperhermitian spaces

2.1 Internal vector spaces A set is called internal if it belongs to an enlargement. A $^*\mathbb{C}$ -vector space F is internal if the set F is internal and if the graphs of its binary operations are internal. A linear function $u : F \rightarrow G$ between internal vector spaces is internal if its graph is internal.

Theorem 1 *Internal ${}^*\mathbb{C}$ vector spaces and internal linear applications make up an abelian category (denoted Ivs).*

Let E be a complex vector space of U and let us assume that binary operations are in U . If $vss_f(E)$ is the set of finite-dimensional subspaces of E , its enlargement ${}^*vss(E)$ is made of *E internal subspaces (and is itself an internal linear space for the enlargement of binary operations of E).

An internal vector space F is said to be hyperfinite-dimensional, if there is a complex vector space E in U such that F is isomorphic (through an internal isomorphism) to an element of ${}^*vss(E)$.

Corollary 2 *The full subcategory of Ivs set made of hyperfinite-dimensional subspaces is a thick subcategory.*

The importance of these subspaces lies in the following proposition :

Proposition 3 *Let H be a complex vector space of U . Since $\mathbb{C} \subset {}^*\mathbb{C}$, *H is also a complex space and the function :*

$$\begin{aligned} f : \quad H &\rightarrow {}^*H \\ h &\mapsto {}^*h \end{aligned}$$

is an injective linear map whose image is included in an internal hyperfinite-dimensional subspace.

2.2 Definition of hyperhermitian spaces and orthonormal hyperbases

The concept of positive-definite sesquilinear form extends easily to hypercomplex vector spaces. We call hyperhermitian space an internal hyperfinite-dimensional vector space provided with an internal positive-definite sesquilinear form. A number of properties are preserved by transfer, when passing from hermitian to hyperhermitian spaces. Let us mention in particular the orthogonal property and moreover the decomposition of a hyperhermitian space into a direct sum of an internal linear subspace and its orthogonal.

Orthonormal hyperbases play a very important role here : a subset B of a hyperhermitian space F is an orthonormal hyperbasis, if B is an orthonormal set and 0 is the only element of F to be orthogonal to every element of B .

Theorem 4 *Let F be a hyperhermitian vector space with scalar product $\langle, \rangle$ and let u be a symmetric internal linear function for $\langle, \rangle$ over F . There exists an orthonormal hyperbasis B of F made of eigenvectors of u .*

Such a hyperbasis B has an important property: it is hyperfinite i.e. it belongs to the enlargement of the set of finite subsets of some set in U_A . We can then define by enlargement the sum operation $\sum$ for a hyperfinite set of indices. This internal operation has by transfer the following characteristic properties:

1. $\sum_{b \in \emptyset} \lambda_b b = 0$
2. $\sum_{b \in B_1 \cup B_2} \lambda_b b + \sum_{b \in B_1 \cap B_2} \lambda_b b = \sum_{b \in B_1} \lambda_b b + \sum_{b \in B_2} \lambda_b b$, for B_1 and B_2 hyperfinite sets.

This leads to:

Proposition 5 *Let B be an orthonormal hyperbasis of a hyperhermitian space F . Every $x \in F$ could be expressed as: $x = \sum_{b \in B} \langle b, x \rangle b$*

Remark 6 *Infinite hyperfinite sets are uncountable.*

2.3 Hull of a hyperhermitian space Let F be a hyperhermitian space with a scalar product $\langle, \rangle$. We define a hypernorm $|||$ (i.e. a norm with hyperreal values) by $|||x||| = \sqrt{\langle x, x \rangle}$.

Let us consider now the set $F_b \subset F$ of bounded hypernorm elements and define over F_b an equivalence relation $\sim$ by : $x \sim y$ iff $x - y$ has an infinitesimal norm. The operations over F are preserved by passing to the quotient space and make of ${}^e F = F / \sim$ a complex vector space. The scalar product $\langle, \rangle$ is also preserved by passing to the quotient and make of ${}^e F$ a prehilbertian space. A classic result (see [Luxemburg Stroyan]) shows that ${}^e F$ is a Hilbert space. For $x \in {}^e F$, we call shadow of x the class of x in ${}^e F$ and denote it also ${}^\circ x$.

Let B be an orthonormal hyperbasis of F . If F is a non finite-dimensional space, set B is uncountable, while for every bounded x , the subset $\{b \in B / \langle b, x \rangle \text{ is not negligible}\}$ is countable. This prevents the extension of the equality $x = \sum_{b \in B} \langle x, b \rangle b$ to the hull of F .

2.4 The Loeb spectral measure Let F be an infinite-dimensional hyperhermitian space and B an orthonormal basis. The set $\mathfrak{T}_0(B)$ of hyperfinite subsets of B is a Boolean algebra where the union of a increasing sequence (B_n) does not belong to the Boolean algebra unless this sequence is stationary. For every $\xi \in F_b$ we define a positive measure on $\mathfrak{T}_0(B)$ by $\nu_\xi(C) = {}^\circ (\sum_{b \in C} |\langle b, \xi \rangle|^2)$. By the fundamental theorem of P. Loeb, this measure extends to the Boolean σ -algebra $\mathfrak{T}(B)$ generated by $\mathfrak{T}_0(B)$ and depends only on the shadow h of ξ .

Theorem 7 *There is a unique mapping ϖ from the Boolean σ -algebra $\mathfrak{T}(B)$ to the set of orthogonal projectors of ${}^e F$ such that for any $h \in {}^e F$ and any $C \in \mathfrak{T}(B)$: $v_h(C) = \langle h, \varpi(C)h \rangle$.*

2.5 Hull of a symmetric internal linear function over a hyperhermitian space Let v be this function. The linear mapping $v - iI$ (where I is the identity function) is bijective and allows us to define the Cayley transform $c_v = (v + iI)(v - iI)^{-1}$ which is a unitary function that becomes, by passing to the quotient, a unitary function ${}^e c_v$ over ${}^e F$. Let ${}^e F_{v,f}$ be the orthogonal of the eigenspace of ${}^e c_v$ associated to the eigenvalue 1.

The graph $\Gamma_v \subset F \times F$ is an internal subset. The shadows of Γ_v bounded elements in ${}^e(F \times F)$ make up a closed subspace G_v . Note here that ${}^e(F \times F)$ could be easily identified with ${}^e F \times {}^e F$.

Proposition 8 *The intersection $G_v \cap {}^e F_{v,f} \times {}^e F_{v,f}$ is the graph of a self-adjoint operator t_v in ${}^e F_{v,f}$.*

Let us define an orthonormal hyperbasis B of F made of v eigenvectors. The mapping $b \mapsto \lambda_b$ (which associates to an eigenvector b its eigenvalue) is an internal hyperreal function. The set $B_{v,f}$ of v eigenvectors with bounded eigenvalues is a borelian set in the sense of 2.4

Proposition 9 *The orthogonal projector over ${}^e F_{v,f}$ is the orthogonal projector associated to the borelian set $B_{v,f}$ by the spectral measure ϖ defined in 2.4*

For any $\lambda \in \mathbb{R}$ the set $\Omega_\lambda = \{b \in B_{v,f} \mid {}^\circ \lambda_b \leq \lambda\}$, is a borelian set. We can thus define $E_\lambda = \varpi(\Omega_\lambda)$ as an orthogonal projector in ${}^e F_{v,f}$.

Proposition 10 *We have $t_v = \int_{-\infty}^{+\infty} \lambda dE_\lambda$, which means that (E_λ) is the spectral family of t_v .*

3 Renormalization of an orthonormal basis

We still consider a hyperhermitian space F and an orthonormal hyperbasis B of F .

3.1 ${}^e F_{B,h}$ Spaces For any element $h \in {}^e F$, $h = {}^\circ x$ and ${}^e F_{B,h}$ denotes the closed subspace generated by $\varpi(C)(h)$, where C evolves in the Borel algebra $\mathfrak{T}(B)$.

ν_h is the unique measure over $\mathfrak{T}(B)$ such that

$$\nu_h(C) = {}^\circ \left(\sum_{b \in C} | \langle b, x \rangle |^2 \right)$$

for every hyperfinite subset C .

Proposition 11 *a. There is a unique isometry $J_{B,h} : {}^eF_{B,h} \rightarrow L^2(B, \nu_h)$ that links $\varpi(C)(h)$ to the characteristic function class of C in $L^2(B, \nu_h)$.*

b. Let z be a bounded element of F and k its shadow in ${}^eF_{B,h}$. The image $\bar{k}$ of k is the class of the function $f_z : b \mapsto {}^\circ \left(\frac{\langle b, z \rangle}{\langle b, x \rangle} \right)$ if this ratio is bounded and 0 otherwise.

c. Let ζ be the measure $\nu_k - |\bar{k}|^2 \nu_h$. ζ and ν_h are mutually singular (and $\nu_k = |\bar{k}|^2 \nu_h + \zeta$ is the Lebesgue decomposition of ν_k with respect to ν_h).

Remark 12 *a. The quantity $\frac{\langle b, z \rangle}{\langle b, x \rangle}$ is in general the ratio of two infinitesimals. We cannot indeed express it through the shadows h and k of x and z .*

b. We will determine $J_{B,h}^{-1}$ in the next paragraph (Theorem 14c).

3.2 Introduction of a second scalar product Let us consider a second internal scalar product $[\cdot, \cdot]$ over the hyperhermitian space F . There exists a symmetric positive definite internal linear function j such that $[x, y] = \langle j(x), j(y) \rangle$. The dual scalar product $\{\cdot, \cdot\}$ is defined by $\{x, y\} = \langle j^{-1}(x), j^{-1}(y) \rangle$ and we have $|\langle x, y \rangle|^2 \leq [x, x] \{y, y\}$ for every x, y .

We now introduce the mapping :

$$\begin{aligned} {}^eF_{[\cdot]} \times {}^eF_{\{\cdot\}} &\rightarrow \mathbb{C} \\ (a, b) &\mapsto \langle a, b \rangle \end{aligned}$$

where : $\langle {}^\circ x_{[\cdot]} | {}^\circ y_{\{\cdot\}} \rangle = {}^\circ \langle x, y \rangle$,

${}^eF_{[\cdot]}$ is the hull of F with a scalar product $[\cdot, \cdot]$,

${}^eF_{\{\cdot\}}$ is the hull of F with a scalar product $\{\cdot, \cdot\}$,

${}^\circ x_{[\cdot]}$ is the shadow of x in ${}^eF_{[\cdot]}$,

${}^\circ y_{\{\cdot\}}$ is the shadow of y in ${}^eF_{\{\cdot\}}$.

Reminder - Let G be a complex vector space. A mapping $f : G \rightarrow \mathbb{C}$ is called antilinear if $f(x + y) = f(x) + \overline{f(y)}$ and $f(\lambda x) = \overline{\lambda} f(x)$. This property is equivalent to have $\overline{f} : x \mapsto \overline{f(x)}$ linear.

A linear function f over a normed space G is bounded, if the linear form $\overline{f}$ is bounded. The norm of f is then defined as the norm of $\overline{f}$. The vector space of bounded antilinear functions from G to $\mathbb{C}$, is called the antilinear dual of G and denoted $\widetilde{G}$.

Proposition 13 *Every linear form f over ${}^eF_{\{\cdot\}}$ can be defined as $y \mapsto \langle x_f, y \rangle$ with a unique $x_f \in {}^eF_{[\cdot]}$. The mapping $f \mapsto x_f$ is an antilinear bijective isometry from the dual of ${}^eF_{\{\cdot\}}$ into ${}^eF_{[\cdot]}$.*

Every antilinear form ϕ over ${}^eF_{[\cdot]}$ can be defined as $x \mapsto \langle x | y_\phi \rangle$ with a unique $y_\phi \in {}^eF_{\{\cdot\}}$. The mapping $y_\phi \mapsto \phi$ is an antilinear bijective isometry from the dual of ${}^eF_{[\cdot]}$ into ${}^eF_{\{\cdot\}}$.

In other words, the sesquilinear form $\langle \cdot | \cdot \rangle$ puts in duality ${}^eF_{[\cdot]}$ and ${}^eF_{\{\cdot\}}$.

In addition we can find an orthonormal hyperbasis Γ of $F, [\cdot]$ which is also orthogonal for $\langle, \rangle$.

The hypersum $\sigma = \sum_{\gamma \in \Gamma} \|\langle \gamma, \gamma \rangle\| = \sum_{\gamma \in \Gamma} \sqrt{\langle \gamma, \gamma \rangle}$ is independent of the choice of Γ basis.

Theorem 14 *Assume σ bounded.*

- a. *The identity mapping from F to F transforms bounded elements for $[\cdot]$ into bounded elements for $\langle, \rangle$ and bounded elements for $\langle, \rangle$ into bounded elements for $\{\cdot\}$. This induces, by passing to the quotient, continuous mappings $: {}^eF_{[\cdot]} \rightarrow {}^eF$ and ${}^eF \rightarrow {}^eF_{\{\cdot\}}$.*
- b. *The complementary of the set $\left\{ b \mid \frac{b}{\langle x, b \rangle} \text{ is bounded for } \{\cdot\} \right\}$ is of null ν_k -measure. Let us define the following function :*

$$\begin{aligned} \phi_x : B &\rightarrow {}^eF_{\{\cdot\}} \\ b &\mapsto \begin{cases} \circ \left(\frac{b}{\langle x, b \rangle} \right) & \text{if } \frac{b}{\langle x, b \rangle} \text{ is bounded} \\ 0 & \text{if not} \end{cases} \end{aligned}$$

- c. *Let $f \in L^2(B, \nu_h)$ and $J_{B,h} : {}^eF_{B,h} \rightarrow L^2(B, \nu_h)$ the isometry defined in proposition 11. The mapping :*

$$\begin{aligned} B &\rightarrow {}^eF_{\{\cdot\}} \\ b &\mapsto f(b)\phi_x(b) \end{aligned}$$

is weakly integrable. For every $q \in {}^eF_h$ we have :

$$\langle \tilde{q}, J_{B,h}^{-1}(f) \rangle = \int_B f(b) \langle q, \phi_x(b) \rangle d\nu_h(b)$$

where $\tilde{q}$ is the image of q by the mapping $: {}^eF_{[\cdot]} \rightarrow {}^eF$ defined in a.

Remark 15 *a. Point c. could be also expressed as : the weak integral $\int_B f(b) \langle q, \phi_x(b) \rangle d\nu_h(b)$ is the image of $J_{B,h}^{-1}(f)$ by the mapping $: {}^eF \rightarrow {}^eF_{\{\cdot\}}$ defined in point a.*

- b. With a possible exception for a countable set of elements b , the scalar product $\langle x, b \rangle$ is infinitesimal and $\frac{b}{\langle x, b \rangle}$ is thus unbounded for $\langle, \rangle$. The introduction of the $\{, \}$ associated norm (renormalization procedure) allows however to extract a finite value.

4 Application to spectral theory

Let G be a complex vector space with a locally convex topology, $\langle, \rangle$ a continuous scalar product over G , and u a continuous linear function over G , symmetric and essentially selfadjoint relative to $\langle, \rangle$. We have the following results :

- a. $\langle u(x), y \rangle = \langle x, u(y) \rangle$ for every x and y in G ,
- b. If $\widehat{G}$ is the completion of G for the scalar product $\langle, \rangle$, the closure of the graph of u in $\widehat{G} \times \widehat{G}$ is the graph of a selfadjoint operator $\bar{u}$.

Note

1. A scalar product $[,]$ over G is nuclear (relative to $\langle, \rangle$) if there is a constant C such that $\sum_{i=1}^n \|b_i\|^2 = \sum_{i=1}^n \langle b_i, b_i \rangle \leq C$ for any finite sequence $b_1, \dots, b_n$ of vectors of G that is orthonormal for $[,]$.
2. $[,]$ is called strongly nuclear (relative to $\langle, \rangle$) if there is an auxiliary nuclear scalar product $[,]_{aux}$ such that $[,]$ is nuclear relative to $[,]_{aux}$.

In what follows we assume a fixed scalar product $[,]$ over G continuous and strongly nuclear relative to $\langle, \rangle$. The existence of such a scalar product is guaranteed if G is a nuclear space.

We introduce the enlargement *G of G and provide it with the internal scalar product obtained by extending the scalar product $\langle, \rangle$ (with the same notation). We then extend u into an internal linear application *u over *G . By proposition 3 there exists a hyperfinite-dimensional internal subspace F of *G that contains the image of G in *G by the mapping $*$.

Let us fix such a subspace F that we assume hyperhermitian for the scalar product $\langle, \rangle$. To simplify notations, we identify G with its image in F .

The canonical injection of G into F induces an isometry from G into eF ; and the closure $\overline{G}$ of G in eF is a completion of G for $\langle, \rangle$.

Let v be a mapping over F that associates to an element x the orthogonal projection of $u(x)$ over F . v is an internal linear application, symmetric for $\langle, \rangle$, that extends u . We can thus apply to v propositions 8 and 9 using same notations :

- Proposition 16** *a. The mapping t_v associated to v in proposition 8 is defined everywhere in G ; and $t_v = u(x)$.*
- b. Let (E_λ) be the spectral family associated to t_v in proposition 10. Every E_λ preserves the closure $\overline{G}$ and the family $(E_{\lambda|\overline{G}})$ of E_λ restrictions to $\overline{G}$ is the spectral family of projectors of the closure $\overline{u}$ of u .*

Remark 17 *Passing from G, u to F, v can be seen as the nonstandard variant of the Ritz projection method.*

4.1 Disintegration of the measure ν_h Let us fix an orthonormal hyperbasis B of eigenvectors of v and denote λ_b the eigenvalue associated to eigenvector b and ϖ the Loeb spectral measure associated to B in theorem 7. Let h be an element of $\overline{G}$. Propositions 9 and 16 show that $\sigma_h(\lambda) = \langle h, E_\lambda h \rangle$ is equal to $\langle h, \varpi(\Omega_\lambda)(b) \rangle$ where Ω_λ is the set of b such that λ_b is bounded and ${}^\circ\lambda_b \leq \lambda$.

Proposition 18 *Let q be a real function defined on B as follows :*

$$q(b) = \begin{cases} {}^\circ\lambda_b & \text{if } \lambda_b \text{ is limited} \\ 0 & \text{otherwise} \end{cases}$$

The Stieljes measure generated by the increasing real function σ_h is the image of measure ν_h by q .

Since a linear raising $j : L^\infty(\mathbb{R}, d\sigma_\lambda) \rightarrow \mathcal{L}^\infty(\mathbb{R}, d\sigma_\lambda)$ (i.e. a linear mapping such as ϕ is the class of $j(\phi)$) exists, with $\mathcal{L}^\infty(\mathbb{R}, d\sigma_\lambda)$ the space of essentially bounded numerical functions, we have the following proposition :

Proposition 19 *We can associate to every hyperfinite subset C of B an element ω_C of the space of essentially bounded numerical functions $\mathcal{L}^\infty(\mathbb{R}, d\sigma_\lambda)$, that meets the following conditions :*

- a. The function ω_C takes only positive real values*
- b. $\omega_\emptyset(\lambda) = 0$ and $\omega_{C_1 \cup C_2}(\lambda) = \omega_{C_1}(\lambda) + \omega_{C_2}(\lambda)$ if C_1 and C_2 are disjoint.*
- c. Every numerical function f integrable with respect to $d\sigma_\lambda$ verifies:*

$$\int_C f(q(b)) d\nu_h(b) = \int_{-\infty}^{+\infty} f(\lambda) \omega_C(\lambda) d\sigma_h(\lambda).$$

Statements a. and b. show that we can extend the mapping $\tau_\lambda : C \mapsto \omega_C(\lambda)$ to a Loeb measure over the Boolean σ -algebra $\mathfrak{T}(B)$.

Proposition 20 a. $\nu_h(C) = \int_{-\infty}^{+\infty} \tau_\lambda(C) d\sigma_h(\lambda)$ for every borelian set C .

b. For σ_h -almost every λ , the measure τ_λ is carried by the set $q^{-1}(\lambda) = \{b \in B \mid \lambda_b \text{ is bounded and } {}^\circ \lambda_b = \lambda\}$

This means that the family (τ_λ) is a disintegration of the measure ν_h with respect to its image by q .

We assumed so far the existence of a scalar product $[\cdot, \cdot]$ that is strongly nuclear over G . This scalar product extends to *G and induces a scalar product on F with hypercomplex values. We will keep denoting it $[\cdot, \cdot]$. We can now introduce the dual scalar product $\{\cdot, \cdot\}$ (see 3.2) :

Lemma 21 a. Let Γ be an orthonormal basis for $[\cdot, \cdot]$ that is also orthogonal for $\langle \cdot, \cdot \rangle$. The hypersum $\sum_{\gamma \in \Gamma} \|\gamma\|$ is bounded – and we can thus apply theorem 14.

b. Let ϕ be an element of F limited for the dual scalar product $\{\cdot, \cdot\}$. For every $g \in G$ the hypercomplex $\langle g, \phi \rangle$ is limited and the function $\phi^G : g \mapsto {}^\circ (\langle g, \phi \rangle)$ is a continuous antilinear form on G .

Consequences of lemma 21

- a. $\eta_b = \frac{b}{\sqrt{\{b, b\}}}$ is of norm 1 for the dual scalar product $\{\cdot, \cdot\}$ and defines therefore the continuous antilinear form η_b^G on G .
- b. Assume again theorem 14 notations. If h is an element of eF and a shadow of an element x , we know that ν_h -almost every $\frac{b}{\langle x, b \rangle}$ is limited for $\{\cdot, \cdot\}$ and we have therefore a continuous antilinear form $\phi_x^G : g \mapsto {}^\circ \left(\frac{\langle g, b \rangle}{\langle x, b \rangle} \right)$.

Consider an element h of G . h is its own shadow. Since

$$\frac{b}{\langle h, b \rangle} = \frac{\frac{b}{\sqrt{\{b, b\}}}}{\langle h, \frac{b}{\sqrt{\{b, b\}}} \rangle}, \quad \frac{b}{\langle h, b \rangle} \text{ is limited for } \{\cdot, \cdot\} \text{ iff } \langle h, \frac{b}{\sqrt{\{b, b\}}} \rangle \text{ is not in-}$$

finitesimal (i.e. if $\eta_b^G(h) \neq 0$). And we have : $\phi_b^G(b)(g) = \frac{\eta_b^G(g)}{\eta_b^G(h)}$.

4.2 Main Theorem Let G'_a be the antilinear dual of G (i.e. the space of continuous antilinear forms on G). G'_a is obviously a complex subspace of the space of complex functions defined on G .

The function u admits an adjoint u^* which is the linear function from G'_a into itself defined by $u^*(\phi)(g) = \phi(u(g))$. We will call eigenfunctional of u any element of G'_a which is an eigenvector of u^* . Those of η_b^G which

are not null are such eigenfunctionals of u (for the eigenvalues $\circ\lambda_b$).

We finally get to the last and main theorem of this work (with the same notations used so far) :

Theorem 22 *Let h be an element of G and B_h the set of $b \in B$ such that $\eta_b^G(h) \neq 0$.*

- a. The set B_h is a borelian subset of B whose complementary is of null σ_h -measure. For $b \in B_h$ the antilinear form η_b^G is an eigenfunctional of u for the eigenvalue $\circ\lambda_b$.*
- b. The family $\left(\phi_h^G(b) = \frac{\eta_b^G}{\eta_b^G(h)}\right), b \in B_h$, is weakly τ_λ -integrable for σ_h -almost every λ .*
- c. For almost every λ the weak integral $\omega_{h,\lambda} = \int_{B_h} \frac{1}{\eta_b^G(h)} \eta_b^G d\tau_\lambda(b)$ is, if not null, an eigenfunctional of u for the value λ .*
- d. Let (E_λ) be the spectral projector family of the closure of u in the completion of G for $<, >$ (see proposition 16). For any $g \in G$ we have $<g, E_\lambda h > = \int_{-\infty}^\lambda \omega_\mu(g) d\sigma_h(\mu)$*

Variant – The antilinear form $g \mapsto <g, E_\lambda h >$ associated to an element $E_\lambda h$ of the completion of G , is the weak integral $\int_{-\infty}^\lambda \omega_\mu d\sigma_h(\mu)$ where for almost every μ the antilinear form ω_μ is, if not null, an eigenfunctional of u for the eigenvalue μ of u .

Remark 23 *The statement c. of the preceding theorem is a nonstandard approach of the integral representation of the spectral projectors of an essentially selfadjoint operator with eigenfunctionals (see [Gelfand Chilov] Tome 3 Chapter 4 n° 4.2 Theorem 1).*

The outcome of this approach is the description of these eigenfunctional ω_t with the eigenvectors of an extension v of the considered operator u in hyperfinite dimension. These eigenvectors do not have an immediate interpretation in the initial space G ; but their normalised forms for the scalar product $\{, \}$ define eigenfunctionals.

We aggregate through integration those eigenfunctionals which are associated to the same eigenvalue; and here again the Loeb's technique comes to make things much easier.

Acknowledgements

I would like to thank Professor François Aribaud for his helpful comments and suggestions. All errors and omissions remain mine.

References

- [Gelfand Chilov] - I.M Gelfand and G.E Chilov - Les distributions
Tome 3 (Equations différentielles) Dunod Paris
1965
- [Goldblatt] - Robert Goldblatt - Lectures on the Hyperreals -
An Introduction to Nonstandard Analysis Graduate
Texts in Mathematics n° 188 - Springer Verlag
1998
- [Jech] - Thomas Jech - Set Theory - Pure and Applied
Mathematics - Academic Press 1978
- [Loeb Wolff] - Peter A. Loeb and Manfred Wolff - Nonstandard
Analysis for the working mathematician Mathe-
matics and its applications vol 510 Kluwer Aca-
demic Publishers 2000
- [Luxemburg Stroyan] - W.A.J Luxemburg and K.D.Stroyan - Introduc-
tion to the Theory of Infinitesimals - Academic
Press 1976